

Brief Summary of Ito Calculus

1. B. Oksendal, Stochastic Differential Equations, Springer 1998.

We have seen SDEs several times now and have a reasonable understanding of the meaning of a process of the form:

$$dX_t = b(X_t)dt + \sigma(X_t)dB_t, \quad t \geq 0$$

$\searrow \mathbb{R}^d$ $\searrow \in \mathbb{R}^{d \times d'}$ $\searrow \mathbb{R}^{d'}$

In this note, we formalize some of the constructions and the calculus associated with this process. In the integral form, the above equation is:

$$X_t = X_0 + \int_0^t b(X_s)ds + \underbrace{\int_0^t \sigma(X_s)dB_s}_{\text{Ito Integral.}}$$

The prob. space we use is denoted by $(\Omega, \mathcal{F}, P)$.

Recall that the **Brownian motion** is an independent increment process, with $B_0 = 0$, and

$$(B_{t+h} - B_t) \sim N(0, h).$$

The term: $\int_0^t b(X_s) ds$ is clear, and is defined


through the usual Riemann sum limit.

$\int_0^t \sigma(X_s) dB_s$ is an Ito Integral. Both $\sigma(X_s)(\omega)$

and $B_s(\omega)$ are random processes.

For a "simple" random process of the form

$$f_t(\omega) = \sum_{j=1}^n e_j(\omega) \cdot \chi_{[t_j, t_{j+1})}(t), \quad \begin{matrix} t_0 = 0 \\ t_n = t \end{matrix}$$



we define

$$\int_0^{\tau} f_t(\omega) dB_s(\omega) = \sum_j e_j(\omega) \cdot (B_{t_{j+1}}(\omega) - B_{t_j}(\omega))$$

In general, we can approximate any "reasonable" function f by simple functions over increasing finer discretization of time, and take a limit in the mean-square sense ($h \rightarrow 0$). Details can be found in [1].

Furthermore, using such a discretization, using B.M. properties described above, we have **Ito Isometry**:

$$E \left[\left(\int_0^t f_s(\omega) dB_s(\omega) \right)^2 \right] = E \left[\int_0^t f_s^2(\omega) dt \right]$$

We also state some of the standard properties of the Ito Integral (these are proved using elementary functions, and taking the limit in $L^2(P)$).
mean square limit

$$1. \quad \int_r^t f_s(\omega) dB_s(\omega) = \int_r^u f_s(\omega) dB_s(\omega) + \int_u^t f_s(\omega) dB_s(\omega)$$

$$2. \quad \int_r^t (c f_s(\omega) + g_s(\omega)) dB_s(\omega) = c \int_r^t f_s(\omega) dB_s(\omega) + \int_r^t g_s(\omega) dB_s(\omega)$$

$$3. \quad E \left[\int_r^t f_s(\omega) dB_s(\omega) \right] = 0$$

4. The Ito integral is a Martingale

Definition : (Filtration) over $(\Omega, \mathcal{F}, \mathbb{P})$

$\{\mathcal{F}_t, t \geq 0\}$ is a family of σ -algebras $\mathcal{F}_t \subseteq \mathcal{F}$,
 s.t. $\forall s < t, \mathcal{F}_s \subseteq \mathcal{F}_t$
 (i.e., an "increasing" sequence of σ -algebras)

Martingale: $\{M_t\}$ is a martingale w.r.t to $\{\mathcal{F}_t\}$ if

(i) M_t is $\mathcal{F}_t$ -measurable,

(ii) $E[|M_t|] < \infty$

(iii) $E[M_t | \mathcal{F}_s] = M_s$ a.s. $\forall s \leq t$.

Let $M_t(\omega) = \int_0^t f_s(\omega) dB_s(\omega)$, $\mathcal{F}_t = \sigma\{B_s, s \leq t\}$
Ito Integral

Then, $M_t(\omega)$ is a martingale w.r.t $\mathcal{F}_t$, and

$$P\left(\sup_{0 \leq t \leq \tau} |M_t| > a\right) \leq \frac{1}{a^2} E\left[\int_0^\tau f_s^2(\omega) ds\right]$$

(proof uses Doob's maximal inequality)

All of above naturally generalizes to d-dimensional B.M;
see [1] for details.

The Ito Formula

(scalar case)

$$dX_t = b_t(X_t) dt + \sigma_t(X_t) dB_t.$$

$Y_t = g_t(X_t)$, $g_t(x)$ is twice cont. differentiable
over $t \times x \in [0, \infty) \times \mathbb{R}$

Then,

$$dY_t = \frac{\partial}{\partial t} g_t(X_t) dt + \frac{\partial}{\partial x} g_t(X_t) dX_t \\ + \frac{1}{2} \frac{\partial^2}{\partial x^2} g_t(X_t) \cdot (dX_t)^2$$

where

$(dX_t)^2$ computed using these rules:

$$dt \cdot dt = dt \cdot dB_t = dB_t \cdot dt = 0$$

$$dB_t \cdot dB_t = dt.$$

The proof of this follows by approximating dt and dB_t by discrete differences $(\Delta t, \Delta B_t = B_{t+\Delta t} - B_t)$

and letting the approximation time-step h go to '0'.

The limit is taken in the $L_2(P)$ sense, and formalizes the calculation we did when we first studied SDEs in (Note 3). The key ideas are to use Taylor's Theorem, and that $(\Delta t)^2 = o(\Delta t)$, $E[(\Delta B_t)^2] \approx \Delta t$, and $E[\Delta B_t] = 0$.

In the vector case, we have the analogous result: $X_t, Y_t \in \mathbb{R}^d$, $g_{t,k}: \mathbb{R}^d \rightarrow \mathbb{R}$,

$$Y_t = g_t(X_t) \quad g_t(\cdot) = \begin{pmatrix} g_{t,1}(\cdot) \\ \vdots \\ g_{t,d}(\cdot) \end{pmatrix}$$

$$dY_k = \frac{\partial g_{t,k}(X_t)}{\partial t} dt + \sum_i \frac{\partial g_{t,k}(X_t)}{\partial x_i} dX_{t,i} + \frac{1}{2} \sum_{i,j} \frac{\partial^2 g_{t,k}(X_t)}{\partial x_i \partial x_j} (dX_{t,i} dX_{t,j})$$

where

$$dB_{t,i} \cdot dB_{t,j} = \delta_{ij} dt, \quad dB_{t,i} \cdot dt = 0, \quad \text{etc.}$$

$\delta_{ij} = \begin{cases} 1 & \text{if } i=j \\ 0 & \text{o.w.} \end{cases}$

Existence / Uniqueness of Solution to SDE:

$$dX_t = b_t(x_t) dt + \sigma_t(x_t) dB_t$$

Suppose $|b_t(x)| + |\sigma_t(x)| \leq C(1 + |x|)$,
 $t \in [0, \tau]$.

and

$$|b_t(x) - b_t(y)| + |\sigma_t(x) - \sigma_t(y)| \leq C'|x - y|,$$

then the SDE has a unique t -continuous (i.e.,
cont. sample path) solution.

Strong Markov Property

$$dX_t = b(x_t) dt + \sigma(x_t) dB_t.$$

$f(\cdot)$ a bounded fn; τ a stopping time
w.r.t. $\{\mathcal{F}_t\}$.

Then, $E_{x_0=x} [f(x_{\tau+h}) | \mathcal{F}_\tau] =$

$$E_{x_0=x_\tau} [f(x_h)]$$

We have already seen the **Generator** $\mathcal{A}$:

$$\mathcal{A}f(x) = \lim_{t \rightarrow 0} \frac{E_{x_0=x} [f(x_t)] - f(x)}{t}, \quad x \in \mathbb{R}^n$$

For: $dX_t = b(X_t)dt + \sigma(X_t)dB_t,$

recall:

$$\mathcal{A}f(x) = \sum_i b_i(x) \frac{\partial f}{\partial x_i} + \frac{1}{2} \sum_{i,j} (\sigma \sigma^\top)_{ij}(x) \frac{\partial^2 f}{\partial x_i \partial x_j}$$

Dynkin's Formula

(aka Fundamental Thm of
calculus for Ito processes)

Let f a square-integrable fn, τ a stopping time

w.r.t. $\mathcal{F}_t$,

$$E_{x_0=x} [f(x_z)] = f(x) + E_{x_0=x} \left[\int_0^z df(x_s) ds \right]$$

Girsanov's Theorem

We next show that if we have two SDEs that are driven by the same Brownian motion (i.e., a coupling) that have different drift terms, then their distributions are "close" to each other.

Before we study this, we first have the following result:

Theorem 1 (Thm 8.6.1 in [1]) Levy's Characterization of Brownian Motion

Let $X_t = (x_{t,1} \dots x_{t,d})^T \in \mathbb{R}^d$ be

a d -dim. continuous stochastic process over $(\Omega, \mathcal{F}, P)$.

Then, $(a) \iff (b)$:

(a) X_t is a Brownian motion wr.t P



(b) (i) X_t is a martingale wr.t. $(\{\mathcal{F}_t\}, Q)$
where $\mathcal{F}_t = \sigma\{X_s, s \leq t\}$

(ii) $(X_{t,i} \cdot X_{t,j}) - \delta_{ij} t$ is a martingale wr.t.
 $(\{\mathcal{F}_t\}, Q)$.

Lemma 1 (8.6.2 in [1]). ν, μ s.t. $\frac{d\nu}{d\mu} = f$.

X an absolutely integrable r.v. over $(\Omega, \mathcal{F}, P)$. $\mathcal{H} \subseteq \mathcal{F}$:

Then, $E_\mu[f \cdot X | \mathcal{H}] = E_\nu[X | \mathcal{H}] E_\mu[f | \mathcal{H}]$.

Theorem 2: (Girsanov's Theorem) (Thm 8.6.4 in [6])

Let $\gamma_t \in \mathbb{R}^d$ be a Ito process of the form:

$$d\gamma_t = \beta_t(\gamma_t) dt + \sigma_t(\gamma_t) dB_t, \quad t \in [0, T]$$

$B_t \in \mathbb{R}^{d'}$ is a Brownian motion, over $(\Omega, \mathcal{F}, \mathbb{P}_T)$,
 $\beta_t(\cdot) : \mathbb{R}^d \rightarrow \mathbb{R}^d$, and $\sigma_t(\cdot) : \mathbb{R}^d \rightarrow \mathbb{R}^{d \times d'}$ (a $d \times d'$ matrix).

Suppose that $\exists u_t(\cdot), \alpha_t(\cdot)$ (both $\mathbb{R}^d \rightarrow \mathbb{R}^d$)

$$\text{st. } \sigma_t(\gamma_t) u_t(\gamma_t) = \beta_t(\gamma_t) - \alpha_t(\gamma_t),$$

and

$$E_{\mathbb{P}_T} \left[e^{\frac{1}{2} \int_0^T u_s^2(\gamma_s) ds} \right] < \infty$$

(Novikov's condition)

$$\text{Let } M_t = e^{\left(- \underbrace{\int_0^t u_s(\gamma_s) dB_s}_{\sum_{i=1}^d u_{s,i}(\gamma_s) dB_{s,i}} - \frac{1}{2} \int_0^t u_s^2(\gamma_s) ds \right)}$$

$$0 \leq t \leq T,$$

and a new measure $\mathbb{Q}_T$ that is absolutely

continuous w.r.t P_T , with M_t being the Radon-Nikodym derivative, i.e.,

$$dQ_T(\omega) = M_T(\omega) dP_T(\omega) \quad \text{over } \mathcal{F}.$$

Then, we have:

$$(A) \quad \hat{B}_t = \int_0^t u_s(\gamma_s) ds + B_t \quad \text{is a B.M. over } (\Omega, \mathcal{F}, Q_T)$$

(B) The process γ_t has the Stochastic Integral

$$\text{representation: } d\gamma_t = \alpha_t(\gamma_t) dt + \sigma_t(\gamma_t) d\hat{B}_t$$

Proof: We prove in two parts.

Step (A): We show that

$$\hat{B}_t = \int_0^t u_s(\gamma_s) ds + B_t \quad \text{is a B.M. over } (\Omega, \mathcal{F}, Q_T)$$

$$\text{i.e., } d\hat{B}_t = u_t(\gamma_t) dt + dB_t$$

Here, to avoid technicalities, we assume $u_t(\cdot)$ is bounded. To show $\hat{B}_t$ is a B.M. over $(\Omega, \mathcal{F}, Q_T)$, we use Levy's char. of B.M. and verify Theorem 1, b(i) and b(ii).

b(i): $\hat{B}_t$ is a martingale w.r.t. Q_T :

$$\text{Let } K_t = M_t \cdot \hat{B}_t, \quad M_t \in \mathbb{R}, \hat{B}_t \in \mathbb{R}^d, K_t \in \mathbb{R}^d.$$

$$\text{i.e., } K_{t,i} = M_t \cdot \hat{B}_{t,i}, \quad i=1,2,\dots,d.$$

where

$$M_t = \exp \left(- \int_0^t u_s(\gamma_s) dB_s - \frac{1}{2} \int_0^t u_s^2(\gamma_s) ds \right)$$

$u dB \equiv \langle u, dB \rangle$ $u^2 = \langle u, u \rangle$

Applications of Ito's Formula: (problems 4.3, 4.4 in [1])

$$\textcircled{a} \quad d(\gamma_t X_t) = \gamma_t dX_t + X_t d\gamma_t + dX_t \cdot d\gamma_t.$$

$$\textcircled{b) } \quad Z_t = e^{-\int_0^t \theta_s(z_s) dB_s - \frac{1}{2} \int_0^t \theta_s^2(z_s) ds}$$

$$\text{Then } dZ_t = -Z_t \cdot \theta_t(z_t) dB_t, \quad \begin{array}{l} Z_t \in \mathbb{R} \\ \theta_t(\cdot) \in \mathbb{R}^d, \\ B_t \in \mathbb{R}^d \end{array}$$

$$\theta_t(\cdot) dB_t \triangleq \sum_{i=1}^d \theta_{t,i}(\cdot) dB_{t,i}$$

$$\therefore dK_{t,i} \stackrel{\textcircled{a)}}{=} M_t d\hat{B}_{t,i} + \hat{B}_{t,i} dM_t + d\hat{B}_{t,i} \cdot dM_t$$

$$= M_t (u_{t,i}(\gamma_t) dt + dB_{t,i})$$

$$+ \hat{B}_{t,i} \underbrace{\left(-M_t \sum_{k=1}^d u_{t,k}(\gamma_t) dB_{t,k} \right)}_{dM_t} \quad \rightarrow \text{from } \textcircled{b)}$$

$$+ (u_{t,i}(\gamma_t) dt + dB_{t,i}) \left(-M_t \sum_{k=1}^d u_{t,k}(\gamma_t) dB_{t,k} \right)$$

$$dt \cdot dB_{t,i} = 0$$

$$dB_{t,i} \cdot dB_{t,j} = \delta_{ij}$$

$$= M_t \left(dB_{t,i} - \hat{B}_{t,i} \sum_{k=1}^d u_{t,k}(\gamma_t) dB_{t,k} \right)$$

$$= M_t (\gamma_t^{(i)} \cdot dB_t), \quad \begin{array}{l} (\gamma_t^{(i)} \in \mathbb{R}^d, B_t \in \mathbb{R}^d) \\ M_t \in \mathbb{R}. \end{array}$$

$$\text{where } \gamma_{t,j}^{(i)} = \begin{cases} -\hat{B}_{t,i} u_{t,j}(\gamma_t) & j \neq i \\ 1 - \hat{B}_{t,i} u_{t,i}(\gamma_t) & j = i \end{cases}$$

i.e.,

$$K_{t,i} = \int_0^t M_s (\gamma_s^{(i)} \cdot dB_s) \text{ is an Ito Integral,}$$

and thus $K_{t,i}$ is a martingale w.r.t. $\mathbb{P}$.

$$E_\lambda[F \cdot X | \mathcal{H}] = E_\nu[X | \mathcal{H}] E_\lambda[P | \mathcal{H}], \quad F = \frac{\partial v}{\partial \lambda}$$

$\therefore$ From Lemma 1, $(\nu \rightarrow Q, \mu \rightarrow P, f \rightarrow M_t, X \rightarrow \hat{B}_t)$

$$E_Q[\hat{B}_{t,i} | \mathcal{F}_s] = \frac{E_P[M_t \hat{B}_{t,i} | \mathcal{F}_s]}{E_P[M_t | \mathcal{F}_s]}$$

$$\text{where } \mathcal{F}_t = \sigma\{B_s, s \leq t\} = \sigma\{\hat{B}_s, s \leq t\}.$$

$$\text{But } E_P[M_t | \mathcal{F}_s] = M_s \quad \left(M_t \text{ is a martingale, see Remark 2} \right)$$

$$E_P[M_t \hat{B}_{t,i} | \mathcal{F}_s] = E_P[K_{t,i} | \mathcal{F}_s] = K_{s,i}$$

$$\therefore E_Q \left[\hat{B}_{t,i} \mid \mathcal{F}_s \right] = \frac{K_{s,i}}{M_s} = \hat{B}_{s,i},$$

i.e., $\{\hat{B}_{t,i}\}$ is a martingale.

Similar proof for b(ii); thus from Levy's char., $\hat{B}_{t,i}$ is a B.M.

Step (B): The process γ_t has the Stochastic Integral representation:

$$d\gamma_t = \alpha_t(\gamma_t) dt + \sigma_t(\gamma_t) d\hat{B}_t$$

By definition,

$$d\gamma_t = \beta_t(\gamma_t) dt + \sigma_t(\gamma_t) dB_t$$

$$= \beta_t(\gamma_t) dt + \sigma_t(\gamma_t) (d\hat{B}_t - u_t(\gamma_t) dt)$$

$$= (\beta_t(\gamma_t) - \sigma_t(\gamma_t) u_t(\gamma_t)) dt + \sigma_t(\gamma_t) d\hat{B}_t$$

$$= \alpha_t(\gamma_t) dt + \sigma_t(\gamma_t) d\hat{B}_t$$

□

Remark 1: Summarizing, we have

$$\begin{aligned} d\gamma_t &= \beta_t(\gamma_t) dt + \sigma_t(\gamma_t) dB_t && \text{BM over } (\Omega, \mathcal{F}, P_T) \\ &= \alpha_t(\gamma_t) dt + \sigma_t(\gamma_t) d\hat{B}_t && \text{BM over } (\Omega, \mathcal{F}, Q_T) \end{aligned}$$

Now, let us define a new process $\tilde{\gamma}_t$ as:

$$d\tilde{\gamma}_t = \alpha_t(\tilde{\gamma}_t) dt + \sigma_t(\tilde{\gamma}_t) dB_t \quad \text{BM over } (\Omega, \mathcal{F}, P_T)$$

Thus, $(\gamma_t, \tilde{\gamma}_t)$ are a pair of processes that are coupled by the same B.M. B_t , but with different drift terms.

Observe that this trajectory γ_t driven by $\hat{B}_t$ (under measure Q_T) is the same in distribution as $\tilde{\gamma}_t$ driven by B_t (under measure P_T).

Remark 2: (From 4.4, Oksendal [1])

$$\text{Suppose } \theta_t = (\theta_{t,1} \dots \theta_{t,d}) \in \mathbb{R}^d$$
$$\text{and } M_t = e^{\int_0^t \theta_s dB_s - \frac{1}{2} \int_0^t \theta_s^2 ds}, \quad t \in [0, \tau]$$

$$\text{where } \theta^2 = \theta \cdot \theta.$$

$$\text{Then (a) } dM_t = M_t (\theta_t \cdot dB_t)$$

(shown using Ito's formula)

$$\text{(b) Suppose } E \left[e^{\frac{1}{2} \int_0^\tau \theta_s^2 ds} \right] < \infty \quad (\text{Novikov condition}).$$

Then M_t is a martingale.