

Time discretization Bounds for DDPM

References:

1. G. Reeves and H. Pfister, Information-theoretic proofs for diffusion sampling, arXiv: 2502.02305, 2025.
2. S. Chen, S. Chewi, J. Li, Y. Li, A. Salim and A. Zhang, Sampling is as easy as learning the score: theory for diffusion models with minimum data assumptions, arXiv 2209.11215, 2022.
3. H. Lee, J. Lu and Y. Tan, Convergence for score based generative modeling with polynomial complexity, arXiv: 2206.06227, 2022.
4. J. Benton, V. de Bortoli, A. Doucet and G. Deligiannidis, Nearly d-linear convergence bounds for diffusion models via stochastic localization, arXiv 2308.03686, 2023.
5. D. Guo, S. Shamai and S. Verdú, Mutual Information and minimum mean-square error in Gaussian channels, IEEE Trans. IT, 2007.

(We closely follow [1] in this note.)

We consider the following dynamic:

$$Z_k = Z_{k-1} + \delta_k f_k(Z_{k-1}) + \sqrt{\delta_k} \tilde{W}_k, \quad k=1, 2, \dots, n$$

①

where $Z_0 = 0$, δ_k , $k=1, 2, \dots, n$, $f_k(\cdot): \mathbb{R}^d \rightarrow \mathbb{R}^d$

$\tilde{W}_k \sim N(0, I_d)$, $k=1, 2, \dots, n$ (iid).

Define the following companion process:

$$\gamma_k = \gamma_{k-1} + \delta_k X_0 + \sqrt{\delta_k} \hat{W}_k, \quad \text{where}$$

②

$X_0 \sim \mu_x$ (ground truth data distribution), $X_0 \perp \{\hat{W}_k\}$

and $\gamma_0 = 0$, $\hat{W}_k \sim N(0, I_d)$

independent

Rewrite γ_k : Let $t_k = (\delta_1 + \delta_2 + \dots + \delta_k)$

Then, $\gamma_k = t_k X_0 + W_k$, $W_k = \sum_{i=1}^k \sqrt{\delta_i} \hat{W}_i$

③

$$\gamma_1 = \delta_1 x_0 + \sqrt{\delta_1} \hat{w}_0 ; \quad \gamma_2 = \gamma_1 + \delta_2 x_0 + \sqrt{\delta_2} \hat{w}_2 \\ = (\delta_1 + \delta_2) x_0 + (\sqrt{\delta_1} \hat{w}_0 + \sqrt{\delta_2} \hat{w}_2)$$

...

$$\text{i.e., } W_k \sim N(0, t_k) \quad \left(\because t_k = \sum_{i=1}^k (\sqrt{\delta_i})^2 \right)$$

$$\text{Further } \text{Cov}(W_k, W_m) = \min\{t_k, t_m\} I_d$$

$$\text{Since } \frac{\gamma_k}{t_k} = x_0 + \frac{W_k}{t_k},$$

$\downarrow$
in $\mathcal{M}_x$

$$(\gamma_k / t_k) \sim \left(\mu_x * N\left(0, \frac{1}{t_k} I_d\right) \right) \quad k=1, 2, \dots, n$$

Theorem 1 [1]: Let $\Delta_x = KL\left(\mu(\gamma_1, \gamma_2, \dots, \gamma_n) \parallel \mu(z_1, z_2, \dots, z_n)\right)$

Note: $\mu(z)$ is the prob. dist. / measure associated with a r.v. z (overloaded notation to denote both dist. / measure)

Assume μ_* has finite second moments, and

$$E \left[\|f_k(\gamma_{k-1})\|^2 \right] < \infty \quad \forall k. \quad \text{Then:}$$

$$\Delta_n \leq \frac{\delta_{\max}}{2} \text{tr}(\text{Cov}(x_0))$$

$$+ \sum_{k=1}^n \frac{\delta_k}{2} E \left[\|f_k(\gamma_{k-1}) - E[x_0 | \gamma_{k-1}] \|^2 \right],$$

where $\delta_{\max} = \max \{ \delta_1, \delta_2, \dots, \delta_K \}$.

Remark: Suppose we have the following:

$$\textcircled{a} \quad f_k(y_{k-1}) = E \left[x_0 | \gamma_{k-1} = y_{k-1} \right]$$

Recall that the annealed score is affine in the conditional expectation. Thus, this selection of $f_k(\cdot)$ corresponds to a DDPM algorithm.

(b) Suppose we wish to sample from a terminal state that $\sim \mu_x \star N(0, \varepsilon I_d)$ for some small $\varepsilon > 0$. Then, suppose that we choose: $\delta_k = \left(\frac{\varepsilon}{n}\right)$, $\forall k=1, 2, \dots, n$.

$$(a) + (b) \Rightarrow \Delta_n \leq \frac{1}{2n\varepsilon} \operatorname{tr}(\operatorname{Cov}(X_0))$$

if $X_0 \sim N(\cdot, \Sigma_d)$,
then $\operatorname{tr}(\operatorname{Cov}(X_0)) \approx d$

(c) By changing f_k s.t. it returns a random value whose moments match $E[X_0^x | Y_{k-1} = y_{k-1}]$, then we can get $O\left(\frac{1}{n^\alpha}\right)$; see [1] for details.

Recall: $Y_k = t_k X_0 + W_k$, $k=1, 2, \dots, n$.
 $\hookrightarrow \sim N(0, t_k I_d)$

Then, the following surprising result holds:

Lemma 1: (a) $\{\gamma_k, k=1, 2, \dots, n\}$ is a discrete time Markov process.

(b) $X_0 \perp\!\!\!\perp (\gamma_1, \gamma_2, \dots, \gamma_{k-1}) \mid \gamma_k$

Proof: Consider the difference sequence:

$$V_k = (\gamma_k - \gamma_{k-1}) = \delta_k X_0 + \sqrt{\delta_k} \widehat{W}_k$$

$\hookrightarrow N(0, I_d)$

Then,

$$q_{V_1, \dots, V_n | X_0} (v_1, \dots, v_n | x_0) = \prod_{i=1}^n \frac{1}{(2\pi\delta_i)^{d/2}} e^{-\frac{1}{2\delta_i} \|v_i - \delta_i x_0\|^2}$$

$\downarrow$
density fn.

$$= \left(\prod_{i=1}^n \phi_i(v_i) \right) e^{-\underbrace{\left\langle \sum_{i=1}^k v_i, x_0 \right\rangle}_{= y_k} - \frac{t_k}{2} \|x_0\|^2}$$

$\longrightarrow$ (4)

where $\phi_i(v_i) = \frac{1}{(2\pi\delta_i)^{d/2}} e^{-\frac{1}{2\delta_i} \|v_i\|^2}$

Aside: Neyman-Fisher Factorization Thm: Let

$f(v_1, \dots, v_n; \alpha_0)$ be a density fn with parameter α_0 . Then $T(\vec{v})$ is a sufficient statistic for α_0 if for any prior dist. over α_0 ,

$$f(\alpha_0 | T(v_1, \dots, v_n)) = f(\alpha_0 | T(v_1, \dots, v_n), v_1, \dots, v_n)$$

$$\text{Equivalently, } I(\alpha_0; T(v_1, \dots, v_n)) = I(\alpha_0; (v_1, \dots, v_n))$$

A statistic $T(\vec{v})$ is sufficient if and only if $\exists$ function $h(\cdot) \geq 0$, $g(\cdot, \cdot) \geq 0$ s.t. $f(\vec{v}; \alpha_0) = h(\vec{v}) g(\alpha_0, T(\vec{v}))$

$$\vec{v} = (v_1, v_2, \dots, v_n)$$

$$\text{Observe that } \gamma_k = \sum_{i=1}^k v_i \quad \left(\begin{array}{l} \text{by telescoping,} \\ \text{and } \gamma_0 = 0 \end{array} \right)$$

is a sufficient statistic for α_0 from $(v_1, \dots, v_n)$ by Neyman-Fisher Thm. Thus, we have:

$$\alpha_0 \perp\!\!\!\perp (v_1, v_2, \dots, v_k) \mid \gamma_k$$

$$\text{i.e., } \alpha_0 \perp\!\!\!\perp (\gamma_1, \gamma_2, \dots, \gamma_{k-1}) \mid \gamma_k$$

(b)

The above fact immediately leads to (a) (please see Lemma 1 in [1]). We will instead prove the M.P. property using an alternate proof, also in [1].

$$\text{Let } B_k = \sqrt{\frac{t_{k+1}}{t_k}} \gamma_k - \sqrt{\frac{t_k}{t_{k+1}}} \gamma_{k+1}$$

→ (4)

$$(\because \gamma_k = t_k x_0 + W_k)$$

$$= \sqrt{\frac{t_{k+1}}{t_k}} W_k - \sqrt{\frac{t_k}{t_{k+1}}} W_{k+1}$$

$$\text{Now, } \text{Cov}(W_k, W_m) = \min\{t_k, t_m\} I_d.$$

$$\begin{aligned} \therefore m > k: \text{Cov}(B_k, W_m) &= \sqrt{\frac{t_{k+1}}{t_k}} \cdot t_k \cdot I_d \\ &\quad - \sqrt{\frac{t_k}{t_{k+1}}} t_{k+1} \cdot I_d \\ &= 0 \end{aligned}$$

$$\therefore B_k \perp (W_{k+1}, W_{k+2}, \dots, W_n).$$

∴ Rewriting (4), we have:

$$\gamma_k = \frac{t_k}{t_{k+1}} \gamma_{k+1} + \sqrt{\frac{t_k}{t_{k+1}}} B_k,$$

with $B_k \sim N(0, \delta_{k+1} I_d)$.

⇒ Reversed process $\{\gamma_n, \gamma_{n-1}, \dots, \gamma_1\}$ is Markov (indep. Gaussian increments).

⇔ (Useful fact!)

Forward process $\{\gamma_1, \gamma_2, \dots, \gamma_n\}$ is Markov.

Ⓐ

□

(Note: The reversed process could be time inhomogeneous even when forward is time homogeneous or vice-versa. However, as long as we show one of these processes is Markovian, the other is also Markovian.)

Lemma 2 Under same assumptions in Thm 1, we have:

$$\Delta_n = \sum_{k=1}^n \frac{\delta_k}{2} E \left[\|X_0 - E[X_0 | Y_{k-1}] \|^2 \right]$$

↗ mean-square estimation error

$$- I(X_0; Y_n) + \sum_{k=1}^n \frac{\delta_k}{2} E \left[\|f_k(Y_{k-1}) - E[X_0 | Y_{k-1}] \|^2 \right]$$

↳ mutual information
↳ can be made to '0' by appropriate choice of f_k .

where $I(X_0; Y_n) = KL(\mu(X, Y_n) \parallel \mu(X)\mu(Y_n))$.

Proof:- $\{Y_k\}$ a M.P. with trans. dist. $p_k(\cdot | \cdot)$
 $\{Z_k\}$ a M.P. with trans dist. $q_k(\cdot | \cdot)$.

By definition, and Markovian property, we have:

$$\Delta_n = KL(\mu(Y_1, \dots, Y_n) \parallel \mu(Z_1, \dots, Z_n))$$

$$= \sum_{k=1}^n E \left[\ln \frac{p_k(Y_k | Y_{k-1})}{q_k(Y_k | Y_{k-1})} \right] \rightarrow \textcircled{5}$$

We now derive expressions for the numerator and denominator in the expectation above.

Recall from ①: $Z_k = Z_{k-1} + \delta_k f_k(Z_{k-1}) + \sqrt{\delta_k} \tilde{W}_k$

$$\therefore q_k(y_k | y_{k-1}) = \phi_k(y_k - y_{k-1} - \delta_k f_k(y_{k-1}))$$

$\hookrightarrow N(0, \delta_k I_d)$

$$\therefore -\ln q_k(y_k | y_{k-1}) = \frac{d}{2} \ln(2\pi \delta_k) + \frac{\|y_k - y_{k-1} - \delta_k f_k(y_{k-1})\|^2}{2\delta_k}$$

$$\text{i.e., } -\ln q_k(y_k | y_{k-1}) = \underbrace{\frac{d}{2} \ln(2\pi \delta_k)}_{= -\frac{d}{2} \ln(2\pi e \delta_k)} - \frac{d}{2}$$

$$= \frac{1}{2\delta_k} \|y_k - y_{k-1} - \delta_k f_k(y_{k-1})\|^2 - \frac{d}{2}$$

Also recall: $\gamma_k = \gamma_{k-1} + \delta_k X_0 + \sqrt{\delta_k} \widehat{W}_k$ (from ②).

$\hookrightarrow N(0, I_d)$

$\therefore$ We have:

$$-E[\ln q_k(\gamma_k | \gamma_{k-1})] = \frac{d}{2} \ln(2\pi e \delta_k)$$

$$= \frac{1}{2\delta_k} E\left[\underbrace{\|\gamma_k - \gamma_{k-1} - \delta_k f_k(\gamma_{k-1})\|^2}_{\text{use ②}}\right] - \frac{d}{2}$$

$$= \frac{1}{2\delta_k} E \left[\| \delta_k X_0 + \sqrt{\delta_k} \hat{W}_k - \delta_k f_k(\gamma_{k-1}) \|^2 \right] - \frac{d}{2}$$

$$= \frac{1}{2\delta_k} \left(E \left[\| \delta_k X_0 - \delta_k f_k(\gamma_{k-1}) \|^2 \right] + \overbrace{E \left[\| \sqrt{\delta_k} \hat{W}_k \|^2 \right]} = \delta_k d \right. \\ \left. + E \left[\langle \delta_k X_0 - \delta_k f_k(\gamma_{k-1}), \sqrt{\delta_k} \hat{W}_k \rangle \right] \right) \\ \underbrace{\hspace{10em}}_{\hat{W}_k \perp (X_0, \gamma_{k-1})} \\ - d/2.$$

$$= \frac{\delta_k}{2} E \left[\| X_0 - f_k(\gamma_{k-1}) \|^2 \right]$$

$$= \frac{\delta_k}{2} E \left[\| X_0 - E[X_0 | \gamma_{k-1}] \|^2 \right]$$

$$+ \frac{\delta_k}{2} E \left[\| E[X_0 | \gamma_{k-1}] - f_k(\gamma_{k-1}) \|^2 \right]$$



principle of orthogonality: $\overset{\text{error vector}}{(X_0 - E[X_0 | \gamma_{k-1}])} \perp \overset{\text{any estimate}}{h(\gamma_{k-1})}$
 for any $h(\cdot)$, because $E[X_0 | \gamma_{k-1}]$ is the minimum mean square estimate of X_0 given γ_{k-1} .

Denominator term of ⑤

So far: $-E[\ln q_k(\gamma_k | \gamma_{k-1})] - \frac{d}{2} \ln(2\pi e \delta_k)$

$$= \frac{\delta_k}{2} E[\|x_0 - E[x_0 | \gamma_{k-1}]\|^2]$$

→ ⑥

$$+ \frac{\delta_k}{2} E[\|E[x_0 | \gamma_{k-1}] - f_k(\gamma_{k-1})\|^2]$$

conditional differential entropy

Defn:

$$h(x|z) = E[\ln p(x|z)] \\ = \iint p(x,y) \ln p(x|y) dx dy$$

Numerator term of ⑤

Next, recall that given (x_0, γ_{k-1}) , γ_k is conditional Gaussian (0 mean, $\delta_k \text{Id}$ covariance).

Thus, $h(\gamma_k | \gamma_{k-1}, x_0) = \frac{d}{2} \ln(2\pi e \delta_k)$

We now have:

$$E[\ln p_k(\gamma_k | \gamma_{k-1})] + \frac{d}{2} \ln(2\pi e \delta_k)$$

$$= -h(\gamma_k | \gamma_{k-1}) + h(\gamma_k | \gamma_{k-1}, x_0)$$

$$I(x; \gamma | z) = h(x | z) - h(x | \gamma, z)$$

$$= I(x_0; \gamma_k | \gamma_{k-1})$$

$$= h(x_0 | \gamma_{k-1}) - h(x_0 | \gamma_{k-1}, \gamma_k)$$

$\gamma_{k-1} - \gamma_k - x_0$
is a p.c from
(b), Lemma 2

$$= (h(x_0) - h(x_0 | \gamma_{k-1})) + (h(x_0) - h(x_0 | \gamma_k))$$

$$= I(x_0; \gamma_k) - I(x_0; \gamma_{k-1}).$$

In summary, we have:

$$E \left[\ln p_k(\gamma_k | \gamma_{k-1}) \right] + \frac{d}{2} \ln(2\pi e \delta_k)$$

→ (6)

$$= I(x_0; \gamma_k) - I(x_0; \gamma_{k-1}).$$

Now, substitute (5), (6) in (4), telescope and
note $I(x_0; \gamma_0) = 0$ □

With these lemmas, we prove Theorem 1. This uses the I-MMSE relationship [5].

Recall note 9a on I-MMSE. Rewriting in integral form:

Suppose $X_0 \sim \mathcal{U}_x$, $N \sim \mathcal{N}(0, I_d)$, $0 \leq a < b$:

Then:

$$\begin{aligned} & I(X_0; \sqrt{b} X_0 + N) - I(X_0; \sqrt{a} X_0 + N) \\ &= \frac{1}{2} \int_a^b M(s) ds \end{aligned}$$

where $M(s) = E \left[\| X_0 - E[X_0 | \sqrt{s} X_0 + N] \|^2 \right]$ ↪ ⑦

$$\text{and } M(0) = \text{tr}(\text{Cov}(X_0)).$$

Fact: $M(s)$ is monotone non-increasing in s .

Proof of Theorem 1:

Recall from Lemma 2 that:

$$\Delta_n = \sum_{k=1}^n \frac{\delta_k}{2} E \left[\|X_0 - E[X_0 | \mathcal{Y}_{k-1}] \|^2 \right] \\ - I(X_0; \mathcal{Y}_n) + \sum_{k=1}^n \frac{\delta_k}{2} E \left[\|F_k(\mathcal{Y}_{k-1}) - E[X_0 | \mathcal{Y}_{k-1}] \|^2 \right]$$

We now show that:

$$\sum_{k=1}^n \frac{\delta_k}{2} E \left[\|X_0 - E[X_0 | \mathcal{Y}_{k-1}] \|^2 \right] - I(X_0; \mathcal{Y}_n) \\ \leq \frac{\delta_{\max}}{2} \text{tr}(\text{Cov}(X_0)),$$

which leads to the Theorem statement.

Before we proceed, we note the following property of Mutual Information in our setting:

$$\mapsto N(0, t_k I_d)$$

$$I(X_0; \mathcal{Y}_k) = I(X_0; t_k X_0 + W_k)$$

from (3),
pp. 3.

$$= I(x_0; \sqrt{t_k} x_0 + N) \quad \xrightarrow{N(0, I_d)}$$

(MI is invariant
to 1-1
transformation)

Then, we have the following:

$$I(x_0; \gamma_k) - I(x_0; \gamma_{k-1}) = \frac{1}{2} \int_{t_{k-1}}^{t_k} M(s) ds, \quad$$

where $t_k = \sum_{i=1}^k \delta_i$ ($\because t_k - t_{k-1} = \delta_k$), and $M(s)$ is monotone non-increasing. Thus,

$$\frac{\delta_k}{2} M(t_k) \leq I(x_0; \gamma_k) - I(x_0; \gamma_{k-1}) \leq \frac{\delta_k}{2} M(t_{k-1})$$

$\hookrightarrow$ (8)


Comment: The above bound is important; it allows us to control the "error" that is accumulated in each time step.

$$\text{Here, } M(t_k) = E[\|x_0 - E[x_0 | \gamma_k]\|^2].$$

Thus,

$$\sum_{k=1}^n \frac{\delta_k}{2} E \left[\|X_0 - E[X_0 | \mathcal{Y}_{k-1}] \|^2 \right] - I(X_0; \mathcal{Y}_n)$$

$$= \sum_{k=1}^n \frac{\delta_k}{2} M(t_{k-1}) - I(X_0; \mathcal{Y}_n)$$

telescope 

$$= \sum_{k=1}^n \left(\frac{\delta_k}{2} M(t_{k-1}) + I(X_0; \mathcal{Y}_{k-1}) - I(X_0; \mathcal{Y}_k) \right)$$

⑧

$$\leq \sum_{k=1}^n \frac{\delta_k}{2} (M(t_{k-1}) - M(t_k))$$

$$\leq \frac{\delta_{\max}}{2} \sum_{k=1}^n (M(t_{k-1}) - M(t_k))$$

$$= \frac{\delta_{\max}}{2} (M(0) - M(t_n))$$

 $= \text{tr}(\text{Cov}(X_0))$

$$\leq \frac{\delta_{\max}}{2} \text{tr}(\text{Cov}(X_0))$$



Comments: There are many other results in [1] that are of interest, including

- ① optimization of step sizes (sec 4.1 in [1])
- ② improved rates via moment matching (4.2 in [1])

along with other connections to literature.
